



Antioxidative and hepatoprotective effects of hydroalcoholic extract of *Artemisia absinthium* L. in rat

Ali Mohammadian¹, Shirin Moradkhani², Sara Ataei³, Tavakol Heidary Shayesteh⁴, Mahsa Sedaghat³, Nejat Kheiripour^{1,5}, Akram Ranjbar^{4*}

¹Student Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

²Department of Pharmacogenosy, School of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran

³Department of Clinical Pharmacy, School of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran

⁴Department of Toxicology and Pharmacology, School of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran

⁵Department of Biochemistry, Hamadan University of Medical Sciences, Hamadan, Iran

ARTICLE INFO

Article Type:
Original Article

Article History:
Received: 2 October 2015
Accepted: 12 November 2015

Keywords:
Artemisia
Oxidative toxic stress
ALT
AST
Rat

ABSTRACT

Introduction: *Artemisia absinthium* L. (AA) is a large, diverse genus of the family Asteraceae. AA has long been used as customary herbal medicine in world for the treatment of gastric pain, cardiac stimulation, improvement of memory and for the restoration of declined mental function. The aim of present study was to evaluate the hepatoprotective effects of AA on some factors reflecting the development of oxidative toxic stress in plasma.

Methods: Twenty male rats were equally divided in to 4 groups (5 rats each). Group I acted as control (received normal saline). Treatment groups were II, III and IV which were given *Artemisia* 10, 50 and 100 mg/kg/day respectively only by gavage for 24 hours. After treatment, blood specimens were collected. Liver enzymes such as aspartate aminotransferase (AST) and alanine aminotransferase (ALT) with total antioxidant power (TAP) and total thiol groups (TTG) concentrations were measured.

Results: Levels of ALT, AST and TTG were decreased in the group II compared to the control (group I). ALT and AST in 50 mg/kg group was observed compared with control group. Also, TTG increased in *Artemisia* 50 mg/kg group compared to control group.

Conclusion: Results suggests that alcoholic extract of *Artemisia* can ameliorate liver toxicity in rats through reducing the serum levels of ALT, AST, and oxidative damage.

Implication for health policy/practice/research/medical education:

Artemisia absinthium is able to ameliorate liver toxicity by reducing oxidative damage and might be beneficial in patients using toxic agents.

Please cite this paper as: Mohammadian A, Moradkhani S, Ataei S, et al. Antioxidative and hepatoprotective effects of hydroalcoholic extract of *Artemisia absinthium* L. in rat. J HerbMed Pharmacol. 2016;5(1):29-32.

Introduction

An imbalance between endogenous antioxidant defense system and reactive oxygen species (ROS) level in the body causes oxidative stress, which is associated with overproduction of free radicals from various liver disorders, such as alcoholic liver damage, non-alcoholic fatty liver disease and drug-induced liver injury (1,2). The liver is an important target of the toxicity of drugs and xenobiotics (3). Recently, natural agents with improved effectiveness and safety profiles as a therapy for liver disease have been extensively sought because the administration of drugs sometimes meets with limited therapeutic success and is

usually associated with serious complications, especially the long-term use (4,5). ROS derived from many sources influence macromolecules in the liver result in hepatotoxicity (6). In general, ROS can be scavenged by detoxifying systems within the body, such as glutathione, glutathione peroxidase, and catalase (7). However, quantities of ROS that overwhelm the capacity of the body's defence system lead to a disturbance of homeostasis in ROS production and antioxidant defence. Which in turn eventually damages biological molecules and key cellular components and processes such as lipid peroxidation, enzyme inactivation, and oxidative DNA damage (8). Dietary antioxidant

*Corresponding author: Akram Ranjbar; Faculty of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran. Tel/Fax: +98 813 8380031, Email: a.ranjbar@umsha.ac.ir

supplementation may help to restore this balance eventually inhibition of hepatotoxicity (9,10). Therefore, plant products or alternative medicines that could limit ROS-mediated injuries are essentially needed to protect aid in the protection the liver from against all types of possible damage (11,12). *Artemisia* is a large, diverse genus of the family Asteraceae. *Artemisia absinthium* (AA), which is also known as 'sweet wormwood' and 'Qinghao' has traditionally been used in traditional Chinese medicine for treatment of fever and chills (13). In particular, *Artemisia* and its derivatives have been used clinically in the treatment of drug-resistant malaria while they were reported to have several bioactive functions including antitumor and anti-inflammatory activities (14-16). In addition, coumarins, flavonoids, and other terpenoids constituents present in AA are also reported to have significant pharmacological activities such as antitumor and antibacterial activities that contribute to the therapeutic effects of the herb (17,18). The aim of the present study was to evaluate the hepatoprotective effects of AA on some factors reflecting associated with the development of oxidative toxic stress in plasma.

Materials and methods

Chemicals and reagents

Dithiobis-(2-nitrobenzoic acid) (DTNB) and Tris base, 2,4,6-tripyridyl-s-triazine (TPTZ) from Merck Chemical Co. (Tehran) and alanine aminotransferase (ALT) & aspartate aminotransferase (AST) kit from Parsazemon Co. (Tehran) were used in this study.

Plant material and preparation of extract

Aerial parts of the plant AA were procured from herbs stores. The taxonomic identity of the plant material was authenticated by Herbarium Unit of the Faculty of Pharmacy, Hamadan University of Medical Sciences, Hamadan, Iran with a deposition specimen holder (No: 220). The dried aerial parts of the plant were milled to a fine powder using an electric blender. The methanol extract was prepared by extracting 50 g of the powdered plant material using soxhlet apparatus (20 hours). Thereafter, the resulting methanol extract was reduced in vacuo (40°C), freeze dried and stored at 4°C until further used. The yield was 11.69% (w/w).

Animals and treatments

20 Wistar rats (200 ± 20 g) were purchased from the Animal Center of Pastor Institute, Wuhan University, Wuhan, China. They were housed on 12 hours light-dark cycle at 25 ± 2°C in relative humidity of 30%-60%, and maintained on a standard pellet diet and water ad libitum. The study received clearance from the Institutional Animal Ethical Committee of the Hamadan University of Medical Sciences, Hamadan, Iran.

The experimental animals were divided into 4 equal groups: Group I acted as control (received nothing).

Treatment groups were II, III and IV which were respectively given *Artemisia* 10, 50 and 100 mg/kg/day by gavage for 24 hours (Group I (control), while group II was given *Artemisia* (10 mg/kg/day). Animals of groups III received only *Artemisia* (50 mg/kg/day). Group IV was given *Artemisia* (100 mg/kg/day) with gavage for 24 hours. After treatment, blood specimens were collected.

Serum biochemical analysis

Blood specimens were collected from the abdominal aorta. After centrifuging at 3000 g for 15 minutes, the serum was separated and stored in -80°C. The serum levels of AST and alanine transaminase ALT were determined by using autoanalyzer.

Determination of antioxidative biomarkers

Assay of total antioxidant power

Total antioxidant power (TAP) was measured by ferric reducing ability of plasma (FRAP) method. This method is based on the ability of plasma in reducing Fe³⁺ to Fe²⁺ in the presence of TPTZ. The reaction of Fe²⁺ and TPTZ gives a complex with a blue color and maximum absorbance in 593 nm (19).

Assay of total thiol groups

To evaluate the plasma total thiol groups (TTG), DTNB was used as a reagent. DTNB reacts with thiol molecules and creates a yellow complex which has good absorbance at 412 nm in spectrophotometer (20).

Statistical analysis

All experiments were done in triplicate and results were reported as mean ± SEM (n=5). The data were analyzed by one-way analysis of variance (ANOVA). Statistically significant effects were further analyzed. Means were also compared using Tukey's multiple range tests. Statistical significance was determined at P<0.05.

Results

Table 1 shows the mean ± SE of variables related to either oxidative stress or liver function in the animals tests. A significant increase (P=0.041) in TTG was observed in the group II as compared to control group (group I). ALT of animals-treated with *Artemisia* 50 mg/kg were significantly (P=0.046) lower than that of control group. Levels of ALT in the group II 50 mg/kg *Artemisia* were significantly (P=0.046) lower than those of the control group (group I). Also, 50 mg/kg *Artemisia*, reduced AST, with respect to the control group (P=0.035). No significant differences were observed in the TAP between the groups.

Discussion

This study was aimed to evaluate the hepatoprotective effects of AA on some factors reflecting the development of oxidative toxic stress in plasma. Administration of hydroalcoholic extract of AA would improved the liver

Table 1. Liver function and antioxidative parameters in animal blood test

Groups n = 5	TAP (umol/ml)	TTG (nmol/ml)	ALT (U/ml)	AST (U/ml)
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Control	2.1 ± 0.02	0.12 ± 0.02	24.5 ± 3.8	52.6 ± 13
Artemisia 10 mg/kg	2.4 ± 0.3	0.14 ± 0.02	22.4 ± 2.1	37.1 ± 8.8
Artemisia 50 mg/kg	2.8 ± 0.4	0.77 ± 0.07 ^a	16.3 ± 1.6 ^a	19.2 ± 2.3 ^a
Artemisia 100 mg/kg	2.6 ± 0.7	0.28 ± 0.11	17.9 ± 2.3	31.4 ± 7.8

Abbreviations: AST, aspartate aminotransferase; ALT, alanine aminotransferase; TAP, total antioxidant power; TTG, total thiol groups.

^a Significantly different from control group at $P < 0.05$.

function and the level of oxidative stress parameters such as TTG in blood. Free radicals are the initiators of a redox reaction cascade, which may lead to changes of the chemical structure of biological macromolecules, such as proteins, lipids and DNA, or trouble of human cell metabolism or even tissue injury (21). In vitro and in vivo studies reported the antioxidant capacity of several species of medicinal plants, acting at cellular level, through cell growth stimulation, membrane potential stabilizing or at molecular level, through ROS scavenging, lipid peroxidation, etc. (22). Also, for the evaluation of liver injury serum concentrations of the most commonly used biochemical markers, ALT and AST were determined. Increased levels of these specific hepatic marker enzymes after chemical or immunological intoxication, compared to control rats, indicated considerable hepato-cellular damage and resulting leakage of cytosolic contents into the systemic circulation (23). AA extract seems to be able to preserve the structural integrity of the hepatocellular membrane, as can be seen with the evident reduction of serum ALT and AST activities of pretreated rats in both experimental models. While previous study showed that 80% aqueous-methanolic extract of *A. absinthium* was effective against CCl₄-injury at the concentration of 500 mg/kg (24), our results indicated improving liver function in 50 mg/kg of AA. Previous studies showed the reduction of serum ALT and AST levels (25,26). We elucidated the probable mechanisms of hepato-protection investigating further the effects of AA on the liver antioxidant status. The liver is a principal organ involved in generation of ROS induced by drugs and toxic chemicals (27). In this study, our results suggested that AA could exert its antioxidant or radical scavenging activities thus preventing the formation ROS. AA administration was shown to put off decrease TTG concentration in experimental animals. AA by directly scavenging the free radicals and improving liver function in rats may increase in the glutathion (GSH) content in the rat liver (13). The previous studies showed antioxidant properties of AA, too (15). In addition, to more clarify possible mechanisms of hepato-protective activity of AA, future studies are required investigating the molecular and cellular mechanisms in oxidative toxic stress pathways.

Authors' contributions

All authors contributed to the conception of the work, conducting the study and approval of the final version of

the manuscript, and agreed for all aspects of the work.

Conflict of interests

The authors declared no competing interests.

Ethical considerations

Ethical issues (including plagiarism, misconduct, data fabrication, falsification, double publication or submission, redundancy) have been completely observed by the authors.

Funding/Support

This study was supported by a grant from Vice Chancellor of Research of Hamadan University of Medical Sciences (Grant No: 9207162136).

References

- Videla LA, Rodrigo R, Orellana M, et al. Oxidative stress-related parameters in the liver of non-alcoholic fatty liver disease patients. *Clin Sci*. 2004;106(3):261-268.
- Kabiri N, Ahangar-Darabi M, Setorki M, Rafieian-Kopaei M. The effect of silymarin on liver injury induced by thioacetamide in rats. *J HerbMed Pharmacol*. 2013;2(2):29-33.
- Zhu R, Wang Y, Zhang L, Guo Q. Oxidative stress and liver disease. *Hepatol Res*. 2012;42(8):741-749.
- Salama SM, Abdulla MA, Al-Rashdi AS, Ismail S, Alkiyumi SS, Golbabapour S. Hepatoprotective effect of ethanolic extract of *Curcuma longa* on thioacetamide induced liver cirrhosis in rats. *BMC Complement Altern Med*. 2013;13(1):56.
- Zhang A, Sun H, Wang X. Recent advances in natural products from plants for treatment of liver diseases. *Eur J Med Chem*. 2013;63:570-577.
- Abdollahi M, Ranjbar A, Shadnia S, Nikfar S, Rezaiee A. Pesticides and oxidative stress: a review. *Med Sci Rev*. 2004;10(6):141-147.
- Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative stress and antioxidant defense. *World Allergy Organ J*. 2012;5(1):9-19.
- Allen R, Tresini M. Oxidative stress and gene regulation. *Free Radic Biol Med*. 2000;28(3):463-499.
- Naik SR, Thakare VN, Patil SR. Protective effect of curcumin on experimentally induced inflammation, hepatotoxicity and cardiotoxicity in rats: evidence

- of its antioxidant property. *Exp Toxicol Pathol.* 2011;63(5):419-431.
10. Mossa AT, Heikal TM, Mohafrash SM, Refaie AA. Antioxidant potential and hepatoprotective activity of origanum majorana leaves extract against oxidative damage and hepatotoxicity induced by pirimiphos-methyl in male mice. *J Appl Sci.* 2015;15(1):69.
 11. Olaleye MT, Rocha BJ. Acetaminophen-induced liver damage in mice: effects of some medicinal plants on the oxidative defense system. *Exp Toxicol Pathol.* 2008;59(5):319-327.
 12. Girish C, Pradhan S. Hepatoprotective activities of picroliv, curcumin, and ellagic acid compared to silymarin on carbon-tetrachloride-induced liver toxicity in mice. *J Pharmacol Pharmacother.* 2012;3(2):149.
 13. Bora KS, Sharma A. Neuroprotective effect of *Artemisia absinthium* L. on focal ischemia and reperfusion-induced cerebral injury. *J Ethnopharmacol.* 2010;129(3):403-409.
 14. Msaada K, Salem N, Bachrouch O, et al. Chemical composition and antioxidant and antimicrobial activities of wormwood (*Artemisia absinthium* L.) essential oils and phenolics. *J Chem.* 2015;3:45-50.
 15. Craciunescu O, Constantin D, Gaspar A, Toma L, Utoiu E, Moldovan L. Evaluation of antioxidant and cytoprotective activities of *Arnica montana* L. and *Artemisia absinthium* L. ethanolic extracts. *Chem Cent J.* 2012;6(1):97.
 16. Ahmad F, Khan RA, Rasheed S. Study of analgesic and anti-inflammatory activity from plant extracts of *Lactuca scariola* and *Artemisia absinthium*. *Journal of Islamic Academy of Sciences.* 1992;5(2):111-114.
 17. Erel SB, Reznicek G, Şenol SG, Yavaşoğlu NÜK, Konyalıoğlu S, Zeybek AU. Antimicrobial and antioxidant properties of *Artemisia* L. species from western anatolia. *Turk J Biol.* 2012;36:75-84.
 18. Singh R, Verma PK, Singh G. Total phenolic, flavonoids and tannin contents in different extracts of *Artemisia absinthium*. *J Intercult Ethnopharmacol.* 2012;1(2):101-4.
 19. Benzie IEF, Strain J. The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power: the FRAP assay. *Anal Biochem.* 1996;239(1):70-6.
 20. Hu M, Dillard C. Plasma SH and GSH measurement. *Methods Enzymol.* 1994;233(385):87.
 21. Halliwell B. Free radicals and antioxidants: updating a personal view. *Nutr Rev.* 2012;70(5):257-265.
 22. Krishnaiah D, Sarbatly R, Nithyanandam R. A review of the antioxidant potential of medicinal plant species. *Food and Bioproducts Processing.* 2011;89(3):217-33.
 23. Poli G. Pathogenesis of liver fibrosis: role of oxidative stress. *Mol Aspects Med.* 2000;21(3):49-98.
 24. Janbaz KH, Gilani AU. Evaluation of the protective potential of *Artemisia maritima* extract on acetaminophen and CCl₄ induced liver damage. *J Ethnopharmacol.* 1995;47(1):43-47.
 25. Gilani AU, Janbaz KH. Preventive and curative effects of *Artemisia absinthium* on acetaminophen and CCl₄ induced hepatotoxicity. *Gen Pharmacol.* 1995;26(2):309-15.
 26. Amat N, Upur H, Blažeković B. In vivo hepatoprotective activity of the aqueous extract of *Artemisia absinthium* L. against chemically and immunologically induced liver injuries in mice. *J Ethnopharmacol.* 2010;131(2):478-484.
 27. Marí M, Morales A, Colell A, García-Ruiz C, Fernandez-Checa JC. Oxidative stress in nonalcoholic fatty liver disease. *Studies on Hepatic Disorders.* Springer; 2015:279-308.